

Betti Numbers of GIT quotients of products of projective planes

Francesca Incensi

November 2009

Abstract

We study the GIT quotients for the diagonal action of the algebraic group $SL_3(\mathbb{C})$ on the n -fold product of $\mathbb{P}^2(\mathbb{C})$: in particular we determine a strategy in order to determine the (intersection) Poincaré polynomial of any quotient variety. In the special case $n = 6$ we determine an explicit formula for the (intersection) Betti numbers of a quotient variety, depending only on the combinatorics of the weights of the polarization $m \in \mathbb{Z}_{>0}^6$.

Introduction

Geometric Invariant Theory gives a method for constructing “quotients” varieties for any linear action of a complex reductive algebraic group G on a projective variety X (see [6] and [1] for the general setting). The quotients depend on the choice of an ample linearized line bundle L : in particular Dolgachev-Hu [2] and Thaddeus [7] proved that only a finite number of GIT quotients can be obtained when L varies and gave a general description of the maps relating the various quotients.

In this paper we restrict our attention to the case when $G = \mathbb{P}SL_3(\mathbb{C})$ acts diagonally on X , the n -fold product of $\mathbb{P}^2(\mathbb{C})$, $\mathbb{P}^2(\mathbb{C})^n$ (that we have already studied in [5]) and describe the topology of an arbitrary quotient variety, both geometric and categorical: as a main result we describe an algorithm to compute the (intersection) Poincaré polynomial of any quotient variety. Moreover in the special case $n = 6$ we obtain an explicit formula for the (intersection) Betti numbers of a quotient variety, depending only on the combinatorics of the weights of the polarization $m \in \mathbb{Z}_{>0}^6$.

The contents of the paper are more precisely as follows.

Section 1 is concerned with the main features of the quotients of $G = \mathbb{P}SL_3(\mathbb{C})$ acting on $X = \mathbb{P}^2(\mathbb{C})^n$: we remind the main features of a categorical quotient $X^{ss}(m)//G$ and we describe the G ample cone $C^G(X)$. At the end we discuss the birational maps $\theta_{\pm k}$ that can relate different quotients.

Section 2 discusses small resolutions: we give conditions for a birational map

θ_{+k} to be a small map, and we discuss the problem of the existence of a small resolution.

In Section 3 we give an algorithm that permits to compute the (intersection) Poincaré polynomial of any quotient. The main key is the decomposition theorem of Beilinson-Bernstein-Deligne.

Section 4 contains the special case $n = 6$: Theorem 4.2 contains an explicit formula for the (intersection) Betti numbers of any quotient: this formula depends only on the combinatorics of the polarization $m \in \mathbb{Z}_{>0}^6$.

1 $G = \mathbb{P}SL_3(\mathbb{C})$ acting on $X = \mathbb{P}^2(\mathbb{C})^n$

First of all remind that $\text{Pic}^G(X) \cong \mathbb{Z}^n$: an ample line bundle L over X is determined by $L = L(m) := L(m_1, \dots, m_n)$, where $m_i \in \mathbb{Z}_{>0}$, $\forall i$ (supp. $m_i \geq m_{i+1}$ for all $i : 1, \dots, n-1$). Then the set of semi-stable points $X^{SS}(m)$ is described by the Hilbert-Mumford numerical criterion; let $x = (x_1, \dots, x_n) \in X$ and $|m| := \sum_{i=1}^n m_i$:

$$x \in X^{SS}(m) \Leftrightarrow \begin{cases} \sum_{k, x_k=y} m_k \leq \frac{|m|}{3} \\ \sum_{j, x_j \in r} m_j \leq 2\frac{|m|}{3} \end{cases} \quad (1)$$

for every point $y \in \mathbb{P}^2(\mathbb{C})$ and for every line $r \subset \mathbb{P}^2(\mathbb{C})$. Moreover x is a stable point, $x \in X^S(m)$, iff the numerical criterion (1) is verified with strict inequalities.

The numerical criterion can be restated as follows: if K and J are subsets of $\{1, \dots, n\}$, then we can associate them with the numbers:

$$\gamma_K^C(m) = |m| - 3 \sum_{k \in K} m_k, \quad \gamma_J^L(m) = 2|m| - 3 \sum_{j \in J} m_j.$$

In particular we have: $\gamma_J^C(m) = -\gamma_{J'}^L(m)$ where $J' = \{1, \dots, n\} \setminus J$.

Now consider $K \subset \{1, \dots, n\}$ with $|K| \geq 2$ and the set V_K^C of configurations $(x_1, \dots, x_n) \in X$ where the points indexed by K are coincident and there are no further coincidences and no non-implied collinearities; in the same way, if $J \subset \{1, \dots, n\}$, $|J| > 2$ consider the set V_J^L of configurations $(x_1, \dots, x_n) \in X$ where the points indexed by J are collinear and there are no further collinearities.

In order to study $X^{SS}(m)$ and $X^S(m)$, it is sufficient to consider the subsets V_K^C and V_J^L : in fact

$$V_K^C \subseteq X^{SS}(m) \Leftrightarrow m_i \leq \frac{|m|}{3} \text{ for all } i \text{ and } \gamma_K^C(m) \geq 0,$$

and

$$V_J^L \subseteq X^{SS}(m) \Leftrightarrow m_i \leq \frac{|m|}{3} \text{ for all } i \text{ and } \gamma_J^L(m) \geq 0,$$

with similar statements for $X^S(m)$ (for more details, see [5]).

Moreover we observe that if $m_i < |m|/3$ for all i , then the number of coincident

points is at most $n - 3$.

Now consider a point

$$\xi \in Z(m) := (X^{SS}(m)//G) \setminus (X^S(m)/G) ;$$

this is the image in $X^{SS}(m)//G$ of different, strictly semi-stable orbits, that all have in their closure a closed, minimal orbit Gx , for a certain configuration x that has $|K|$ coincident points, and the others $n - |K|$ collinear; by the numerical criterion, we get $\gamma_K^C(m) = 0$ and $\gamma_{K'}^L(m) = 0$, where K indicates the coincident points, while $K' = \{1, \dots\} \setminus K$ indicates the collinear ones. In this case $\phi(\overline{V_K^C} \cup \overline{V_{K'}^L})$ has dimension $n - |K| - 3$, where ϕ is the projection map to the quotient: $\phi : X^{SS}(m) \rightarrow X^{SS}(m)//G$.

Some quotients are particularly easy to compute: let's consider two examples.

Example 1.1. Consider a polarization $m = (s, s, s, s, 1, 1, \dots, 1, 1)$ such that

$$\begin{cases} 0 < s < \frac{1}{3}(4s + n - 4) \\ \frac{1}{3}(4s + n - 4) < 2s < \frac{2}{3}(4s + n - 4) & \Rightarrow s > 2(n - 4) . \\ 3s > \frac{2}{3}(4s + n - 4) \end{cases}$$

In this case the quotient $X^{SS}(m)//G$ is isomorphic to the product of $(n - 4)$ copies of $\mathbb{P}^2(\mathbb{C})$.

Example 1.2. Consider the polarization $m = (s, s, 1, 1, \dots, 1)$ such that

$$\begin{cases} 0 < s < \frac{1}{3}(2s + n - 2) \\ \frac{1}{3}(2s + n - 2) < 2s < \frac{2}{3}(2s + n - 2) & \Rightarrow n - \frac{7}{2} < s < n - 2 . \\ n - 3 < \frac{1}{3}(2s + n - 2) \end{cases}$$

In this case the quotient $X^{SS}(m)//G$ is isomorphic to the product of $\mathbb{P}^{n-4}(\mathbb{C}) \times \mathbb{P}^{n-4}(\mathbb{C})$.

Dolgachev and Hu proved in [2] that varying the line bundle L only a finite number of different quotients can be obtained: the space that parametrizes these quotients is the G -ample cone $C^G(X)$, the convex cone in $NS^G(X) \otimes \mathbb{R}$ spanned by ample G -linearized line bundles L with $X_L^{SS} \neq \emptyset$, where $NS^G(X)$ is the (Néron-Severi) group of G -line bundles modulo homological equivalence. This cone is subdivided in walls and chambers: a polarization L lies on a wall if and only if $X_L^S \subsetneq X_L^{SS}$. A chamber is a connected component of the complement of the union of walls: all the polarizations in a chamber define the same set of stable points.

In our case

$$C^G(X) = \{(m_1, \dots, m_n; \lambda) \in \mathbb{R}^n \times \mathbb{R}_+ : \sum_{i=1}^n m_i = 3\lambda, 0 \leq m_i \leq \lambda, i = 1, \dots, n\}.$$

Moreover $X^S(m) \subsetneq X^{SS}(m)$ if and only if there exists a subset $K \subset \{1, \dots, n\}$ such that

$$\sum_{i \in K} m_i = \frac{|m|}{3};$$

this is equivalent to the condition that $L(m)$ belongs to the hyperplane

$$H_{C_K, L_{K'}} = \left\{ (m; \frac{|m|}{3}) \in C^G(X) : \sum_{i \in K} m_i = \frac{|m|}{3}, \sum_{j \in K'} m_j = \frac{2}{3}|m|, K \subset \{1, \dots, n\} \right\}.$$

This is a codimension-1 wall.

Now we want to introduce the birational maps that may be constructed between two quotients: they will be crucial in determining the Betti numbers of the quotients.

Let m be a polarization such that 3 divides $|m|$ and $X^S(m) \neq \emptyset, X^S(m) \subsetneq X^{SS}(m)$; let us consider “variations” of m as follows:

$$\widehat{m} = m \pm (0, \dots, 0, \underbrace{1}_k, 0, \dots, 0).$$

Then applying the numerical criterion we get $X^{SS}(m) \subset X^S(\widehat{m}) = X^{SS}(\widehat{m}) \subset X^{SS}(m)$: these inclusions induce a morphism

$$\theta_{\pm k} : X^S(\widehat{m})/G \longrightarrow X^{SS}(m)/G, \quad (2)$$

which is an isomorphism over $X^S(m)/G$, while over $Z(m)$ is a contraction of subvarieties. Moreover if the quotient $X^{SS}(m)/G$ is singular (possible for $n \geq 6$), then $\theta_{\pm k}$ is a resolution of singularities: from now on we assume $n \geq 6$.

Consider a point $\xi \in Z(m)$: for what we have already observed, ξ is determined by a closed, minimal orbit Gx for a certain configuration x that has $|K|$ coincident points and the other $n - |K|$ collinear ($\gamma_K^C(m) = \gamma_{K'}^L(m) = 0$).

Now we want to calculate d , the dimension of $\theta_{\pm k}^{-1}(\xi)$: by the numerical criterion, only one between V_K^C and $V_{K'}^L$ is included in $X^S(\widehat{m})$.

Dealing with an elementary transformation of “plus” type ($\widehat{m} \xrightarrow{+1_k} m$), then

- if $k \in K \Rightarrow$

$$\theta_{+k}^{-1}(\xi) \cong \mathbb{P}^{n-|K|-3}(\mathbb{C}), \quad d = n - |K| - 3; \quad (3)$$

- if $k \in K' \Rightarrow$

$$\theta_{+k}^{-1}(\xi) \cong \mathbb{P}^{2|K|-3}(\mathbb{C}), \quad d = 2|K| - 3. \quad (4)$$

Similarly, with an elementary transformation of “minus” type ($\widehat{m} \xrightarrow{-1_k} m$), we have

$$k \in K \Rightarrow d = 2|K| - 3; \quad k \in K' \Rightarrow d = n - |K| - 3. \quad (5)$$

From now on we will consider only elementary transformation of “plus” type, θ_{+k} ; in particular we want to study the properties of θ_{+k} : when θ_{+k} is a blow-up map? When a small map?

2 Small resolutions

Definition 2.1. *A proper surjective algebraic map $f : Y_1 \rightarrow Y_2$ between irreducible complex N -dimensional algebraic varieties is small if Y_1 is nonsingular and, for all $r > 0$,*

$$\text{codim}_{\mathbb{C}}\{y \in Y_2 \mid \dim_{\mathbb{C}} f^{-1}(y) \geq r\} > 2r.$$

A small resolution is a resolution of singularities which is a small map.

Small maps are particularly relevant, because they preserve intersection homology.

Proposition 2.2. *Let $X^{SS}(m)//G$ be a quotient such that m lies on a 1 codimension wall of $C^G(X)$ and consider a variation of the weights $\widehat{m}$; then the birational map*

$$\theta_{+k} : X^S(\widehat{m})/G \rightarrow X^{SS}(m)//G,$$

is a small resolution if for each $\xi \in Z(m)$, determined by a minimal closed orbit in $\overline{V}_K^G \cup \overline{V}_{K'}^L$ ($2 \leq |K| \leq n - 3$),

$$\dim(\theta_{+k}^{-1}(\xi)) = \begin{cases} 2|K| - 3 & \text{if } 2 \leq |K| < \frac{n+1}{3}, \\ n - |K| - 3 & \text{if } \frac{n-1}{3} < |K| \leq n - 3. \end{cases} \quad (6)$$

Remark 2.3. *Studying the subdivision in chambers and walls of the G -ample cone $C^G(X)$, the existence of a small resolution appears quite natural: consider a polarization m that determines a singular quotient $X^{SS}(m)//G$ and lies on a codimension-1 wall: each singularity of the quotient, determined by a certain $\phi(\overline{V}_K^G \cup \overline{V}_{K'}^L)$, can be solved by the birational morphisms*

$$\theta_{+k} : X^S(m_+)/G \rightarrow X^{SS}(m)//G \quad \text{and} \quad \theta_{-k} : X^S(m_-)/G \rightarrow X^{SS}(m)//G$$

where the polarizations m_+ and m_- lie on opposite chambers. In order to have a small resolution we have to choose the “right side” of each wall.

Proof. The demonstration is based on the definition of *small map*: in order to verify the definition in our case, we have to consider the codimension of the sets

$$\{\xi \in Z(m) \mid \dim(\theta_{+k}^{-1}(\xi)) \geq r\}$$

for $r > 0$. We have to distinguish two different cases:

1. if $\dim(\theta_{+k}^{-1}(\xi)) = 2|K| - 3$, then

$$\begin{aligned} \text{codim} \{ \xi \in Z(m) \mid \dim(\theta_{+k}^{-1}(\xi)) \geq 2|K| - 3 \} &> 2(2|K| - 3) \\ 2(n - 4) - (n - |K| - 3) &> 4|K| - 6 \\ 3|K| &< n + 1. \end{aligned}$$

2. if $\dim(\theta_{+k}^{-1}(\xi)) = n - |K| - 3$, then

$$\begin{aligned} \text{codim} \{ \xi \in Z(m) \mid \dim(\theta_{+k}^{-1}(\xi)) \geq n - |K| - 3 \} &> 2(n - |K| - 3) \\ 2(n - 4) - (n - |K| - 3) &> 2n - 2|K| - 6 \\ 3|K| &> n - 1. \end{aligned}$$

In other words $\theta_{+k} : X^S(\widehat{m}) \rightarrow X^{SS}(m)//G$ is a small map if for every $\xi \in Z(m)$, $\theta_{+k}^{-1}(\xi)$ has minimum dimension between $2|K| - 3$ and $n - |K| - 3$. ■

Remark 2.4. Combining the previous result and formulas (3) and (4), we can restate Proposition 2.2 as follows: in the same hypothesis of 2.2, the birational map θ_{+k} is a small resolution if for each $\xi \in Z(m)$, determined by a minimal closed orbit in $\overline{V}_K^C \cup \overline{V}_{K'}^L$ ($2 \leq |K| \leq n - 3$),

$$\begin{cases} k \in K' \\ 2 \leq |K| < \frac{n+1}{3}, \end{cases} \quad \text{or} \quad \begin{cases} k \in K \\ \frac{n-1}{3} < |K| \leq n - 3. \end{cases} \quad (7)$$

Example 2.5. Let us consider an example: $n = 7, m = (9, 4, 4, 4, 4, 1), |m| = 30$; this polarization lies on the codimension-1 wall $H_{C_{17}, L_{23456}}$. The map $\theta_{+6} : X^S(9, 4, 4, 4, 4, 3, 1)/G \rightarrow X^{SS}(m)//G$ is a small map, while $\theta_{+1} : X^S(8, 4, 4, 4, 4, 4, 1)/G \rightarrow X^{SS}(m)//G$ is not a small map.

As an immediate consequence we have $IH_{\bullet}(X^{SS}(m)//G)$ is isomorphic to $H_{\bullet}(X^S(\widehat{m}))$ if θ_{+k} is a small map.

In general we have the following stronger result regarding the existence of small resolutions whose proof is the same of [4] Theorem 2.5, in the case of a torus action; in fact using the Gelfand-MacPherson correspondence, the moduli space $(\mathbb{P}^2(\mathbb{C})^{SS})(m)//SL_3(\mathbb{C})$ can be identified with a quotient in the Grassmannian $\text{Gr}(3, \mathbb{C}^n)$ acted on by the torus $(\mathbb{C}^*)^{n-1}$.

Theorem 2.6. For every singular quotient $X^{SS}(m)//G$, there exists a polarization $\widehat{m}$ such that $\widehat{m}$ lies on a chamber close enough to m and $\theta : X^S(\widehat{m})/G \rightarrow X^{SS}(m)//G$ is a small resolution.

Remark 2.7. We have already proved the result for m lying on a codimension-1 wall; for higher codimension cases, the proof is based on the observation that a codimension- N wall is the intersection of N codimension-1 walls H_i and for each wall H_i ($i = 1, \dots, N$) $C^G(X) \setminus H_i$ has two connected components lying in two sides of H_i ; one of these components, named $C_{i,s}$, defines a small resolution according to the previous result. Then the proof examines each wall, detects the right component $C_{i,s}$ and studies their intersection for $i = 1, \dots, N$.

Example 2.8. Consider $n = 8$ and the polarization $m = (19, 16, 7, 7, 6, 2, 2, 1)$: it lies on a codimension-3 wall, given by

$$H_{C_{18}, L_{234567}} \cap H_{C_{267}, L_{13458}} \cap H_{C_{345}, L_{12678}} .$$

A polarization $\hat{m}$ such that $\theta : X^S(\hat{m})/G \rightarrow X^{SS}(m)//G$ is a small map is $\hat{m} = (19, 16, 7, 6, 6, 2, 1, 1)$: in fact $X^S(m) \subset X^S(\hat{m}) \subset X^{SS}(m)$ and $\theta^{-1}(\xi)$ has the correct dimension for each $\xi \in Z(m)$. The polarization $\hat{m}$ is obtained studying the three codimension-1 walls in order to detect the right components and their intersection.

3 Poincaré polynomial

Let us consider a quotient $Y = Y(m) = X^{SS}(m)//G$ and define the intersection Betti numbers of $X^{SS}(m)//G$ as

$$ib_i(Y) := \dim IH_i(Y, \mathbb{Q}), \quad i : 0, \dots, 4(n-4) .$$

If Y is non-singular, this definition coincides with the classical one. In what follows we will use

$$P(Y) = \sum_i t^i \dim(H_i(Y))$$

to denote the ordinary Poincaré polynomial of a variety Y and use

$$IP(Y) = \sum_i t^i \cdot ib_i(Y)$$

to denote the intersection Poincaré polynomial of Y .

We have already stressed that small maps preserve intersection homology: if $\theta_{\pm k} : X^S(\hat{m})/G \rightarrow X^{SS}(m)//G$ is a small map, then $H_\bullet(X^S(\hat{m})/G) \cong IH_\bullet(X^{SS}(m)//G)$, and as a consequence the Betti numbers and the Poincaré polynomial are preserved.

In the previous section we have determined the conditions for a θ_{+k} map to be small, but in the general case θ_{+k} is not small. In this way in order to compute the Betti numbers for a general quotient it is necessary to use the decomposition theorem of Beilinson-Bernstein-Deligne (in particular we will consider the simplified version of [4]):

Theorem 3.1. *Let $f : X \rightarrow Y$ be a projective algebraic map and X be a nonsingular variety. Then there exists*

1. a stratification $Y = \bigcup_\alpha Y_\alpha$ of Y
2. a list of enriched strata $E_\beta = (Y_\beta, L_\beta)$ where Y_β is a stratum of Y and L_β is a local system over Y_β ; moreover assume that every local system L_β is trivial.

Then there exists a collection of polynomials ψ_β for all strata such that,

$$P(X) = \sum_{\beta} IP(\overline{Y}_\beta) \cdot \psi_\beta,$$

and for a point $y \in Y$

$$IP(f^{-1}(y)) = \sum_{\beta} IP_y(\overline{Y}_\beta) \cdot \psi_\beta.$$

Consider a polarization m on a codimension-1 wall $H_{C_K, L_{K'}}$, and $\widehat{m}, \overline{m}$ two polarizations close enough to m lying in different sides of the wall $H_{C_K, L_{K'}}$:

$$\begin{array}{ccc} X^S(\widehat{m})/G & & X^S(\overline{m})/G \\ & \searrow \widehat{\theta} & \overline{\theta} \swarrow \\ & X^{SS}(m)//G & \end{array}$$

Consider the map $\widehat{\theta}: X^S(\widehat{m})/G \rightarrow X^{SS}(m)//G$: suppose that for $\xi \in Z(m)$, $\widehat{\theta}^{-1}(\xi) \cong \mathbb{P}^{n-|K|-3}(\mathbb{C})$. Now apply the decomposition theorem to the map $\widehat{\theta}$, where the stratification of $X^{SS}(m)//G$ is given by

$$X^{SS}(m)//G = X^S(m)/G \cup Z(m).$$

Then there exist two polynomials ψ_0 and φ_1 such that

$$P(X^S(\widehat{m})/G) = \psi_0 IP(X^{SS}(m)//G) + \psi_1 IP(Z(m)).$$

These two polynomials may be determined:

$$\psi_0 = 1, \quad \psi_1 = 1 + t^2 + \dots + t^{2(n-|K|-3)} - IP_\xi(X^{SS}(m)//G),$$

where ξ is a point in $Z(m)$.

Then if ξ is any point in $Z(m)$, we have that $P(X^S(\widehat{m})/G)$ is equal to

$$IP(X^{SS}(m)//G) + \left(1 + \dots + t^{2(n-|K|-3)} - IP_\xi(X^{SS}(m)//G)\right) IP(Z(m)). \quad (8)$$

In the same way, study the map $\overline{\theta}: X^S(\overline{m})/G \rightarrow X^{SS}(m)//G$: suppose that for $\xi \in Z(m)$, $\overline{\theta}^{-1}(\xi) \cong \mathbb{P}^{2|K|-3}(\mathbb{C})$.

Then if ξ is any point in $Z(m)$, we have that $P(X^S(\overline{m})/G)$ is equal to

$$IP(X^{SS}(m)//G) + \left(1 + \dots + t^{2(2|K|-3)} - IP_\xi(X^{SS}(m)//G)\right) IP(Z(m)). \quad (9)$$

Subtracting (8) and (9), $P(X^S(\widehat{m})/G) - P(X^S(\overline{m})/G)$ is equal to

$$P(X^S(\widehat{m})/G) - P(X^S(\overline{m})/G) = \varepsilon(H_{C_K, L_{K'}}) Q_t(H_{C_K, L_{K'}}) IP(Z(m)),$$

where $\varepsilon(H_{C_K, L_{K'}})$ and $Q_t(H_{C_K, L_{K'}})$ are defined by

$$\varepsilon(H_{C_K, L_{K'}}) = \begin{cases} 1 & \text{if } n - |K| - 3 > 2|K| - 3 \Rightarrow |K| < \frac{n}{3} \\ -1 & \text{if } n - |K| - 3 < 2|K| - 3 \Rightarrow |K| > \frac{n}{3} \\ 0 & \text{if } n - |K| - 3 = 2|K| - 3 \Rightarrow |K| = \frac{n}{3} \end{cases}$$

$$Q_t(H_{C_K, L_{K'}}) = \begin{cases} t^{2(2|K|-3)+2} + \dots + t^{2(n-|K|-3)} & \text{if } |K| < \frac{n}{3} \\ t^{2(n-|K|-3)+2} + \dots + t^{2(2|K|-3)} & \text{if } |K| > \frac{n}{3} \\ 0 & \text{if } |K| = \frac{n}{3} \end{cases}$$

We can also write

$$P(X^S(\widehat{m})/G) = P(X^S(\overline{m})/G) + \varepsilon(H_{C_K, L_{K'}})Q_t(H_{C_K, L_{K'}})IP(Z(m)). \quad (10)$$

The previous formula tells us how the Poincaré polynomial and the Betti numbers vary, when we cross a codimension-1 wall $H_{C_K, L_{K'}}$; in particular let study $IP(Z(m))$.

The locus $Z(m)$ is equal to $X^{SS}(m)//G \setminus X^S(m)/G$ and it is determined by

$$\phi(\overline{V_K^C} \cup \overline{V_{K'}^L}) \cong \left(\mathbb{P}^1(\mathbb{C})^{n-|K|}(m') \right)^{SS} // SL_2(\mathbb{C}),$$

where m' is obtained from m , by deleting all those weights m_i with $i \in K$. For the Hilbert-Mumford numerical criterion, the open set of stable points for m' is equal to the open set of semi-stable points for m' , i.e. the categorical quotient $(\mathbb{P}^1(\mathbb{C})^{n-|K|}(m'))^{SS} // SL_2(\mathbb{C})$ is also geometric.

Now the Poincaré polynomials for this kind of quotient are well known (see [3] for details):

$$P(Z(m)) = \frac{1}{1-t^2} \sum_{J \in S_{n-|K|}} \left(t^{2|J|} - t^{2(n-|K|-|J|-2)} \right). \quad (11)$$

where $S_{n-|K|} = \{J \subset \{1, 2, \dots, n-|K|\} : m'_{n-|K|} + \sum_{j \in J} m'_j < \sum_{i \notin J} m'_i\}$.

Example 3.2. In Example 2.5 we have introduced the polarization $m = (9, 4, 4, 4, 4, 4, 1)$ that lies on the codimension-1 wall $H_{C_{17}, L_{23456}}$.

We have $Z(m) \cong (\mathbb{P}^1(\mathbb{C})^5(4, 4, 4, 4, 4))^{SS} // SL_2(\mathbb{C})$ and

$$S_5 = \{\emptyset, \{1\}, \{2\}, \{3\}, \{4\}\}.$$

$$\Rightarrow P(Z(m)) = \frac{1}{1-t^2}(1-t^6+4t^2-4t^4) = 1+5t^2+t^4.$$

In fact $Z(m)$ is isomorphic to the blow-up of $\mathbb{P}^2(\mathbb{C})$ in four points.

Now collecting all the results, we can calculate the Poincaré polynomial of any quotient $X^{SS}(m)//G$; this is the strategy:

- 1) If m lies on a face of $C^G(X)$ and there exists one or more weights m_i such that $m_i = 0$, then the quotient has lower dimension than expected: study the polarization m' obtained from m by deleting all the zero weights, and follow the next steps;

- 2) If m lies on a face of $C^G(X)$ and there exists one weight m_i such that $m_i = \frac{|m|}{3}$, then the quotient degenerates to $(\mathbb{P}^1(\mathbb{C})^{n-1}(m'))^{SS} // SL_2(\mathbb{C})$, where m' is obtained from m by deleting m_i ; if m' determines a categorical quotient that is also geometric use the Hausmann-Knutson formula, otherwise apply the decomposition theorem to this case;
- 3) If m lies on a face of $C^G(X)$ and there exists two weights m_i, m_j such that $m_i = m_j = \frac{|m|}{3}$, then the quotient degenerates to a point;
- 4) If m lies on a wall (its codimension is not relevant), then determine a small resolution $X^S(\hat{m})/G$ of $X^{SS}(m)/G$: the Poincaré polynomial does not change; now follow 5) for the polarization $\hat{m}$;
- 5) If m lies in a chamber, then construct a path γ in $C^G(X)$ that goes from a polarization $\tilde{m}$ whose quotient is well-known (see Examples 1.1 and 1.2) to m , and γ meets only codimension-1 walls. Then if γ crosses the codimension-1 walls $H_{C_{K_1}, L_{K'_1}}, \dots, H_{C_{K_N}, L_{K'_N}}$,

$$\gamma : m = \tilde{m} = m_0 \xrightarrow{H_{C_{K_1}, L_{K'_1}}} m_1 \xrightarrow{H_{C_{K_2}, L_{K'_2}}} m_2 \rightarrow \dots \xrightarrow{H_{C_{K_N}, L_{K'_N}}} m_N = m$$

study each crossing and apply result (10):

$$P(X^S(m_{j+1})/G) = P(X^S(m_j)/G) + \varepsilon(H_{C_{K_i}, L_{K'_i}}) Q_t(H_{C_{K_i}, L_{K'_i}}) IP(Z(m_{j,j+1})),$$

where $m_{j,j+1}$ indicates the polarization that lies on the wall $H_{C_{K_{j+1}}, L_{K'_{j+1}}}$ and “connects” m_j and m_{j+1} . The Poincaré polynomial of each $Z(m_{j,j+1})$ can be computed using formula (11).

4 A special case: $n = 6$

Let's study the particular case $n = 6$: it is really interesting because in this case every map θ_{+k} is the eventual composition of a small map and a blow-up map.

First of all remind that the number of chambers in which the G -ample cone $C^G(X)$ is divided is less than or equal to 38 (see [5] for details).

For quotients $X^{SS}(m)/G$ such that $X^S(m) \subsetneq X^{SS}(m)$, it is not so easy to give an upper bound to the number of quotients, but we can get some important informations about their structure: the following result ([5]) classifies the different types of points that may appear in $Z(m)$:

Theorem 4.1. *Let $X = \mathbb{P}^2(\mathbb{C})^6$ and $m \in \mathbb{Z}_{>0}^6$ a polarization with $3 \mid |m|$ and $m_i < |m|/3 \forall i$: if*

1. *there are two different indexes i, j s.t. $m_i + m_j = |m|/3$, then the quotient includes a curve $C_{ij} \cong \mathbb{P}^1(\mathbb{C})$, that corresponds to strictly semi-stable*

orbits s.t. $x_i = x_j$ or x_h, x_k, x_l, x_n collinear. In particular points ξ of C_{ij} are singular: locally, the variety $(X^{SS}(m)//G, \xi)$ is isomorphic to the toric variety

$$\mathbb{C}[T_1, T_2, T_3, T_4, T_5]/(T_1T_4 - T_2T_3).$$

2. there is a “partition” of m such that $m_i + m_j = m_h + m_l = m_k + m_n$, then the quotient includes three curves $C_{ij}, C_{hl}, C_{kn} \cong \mathbb{P}^1(\mathbb{C})$, that have a common point $O_{ij,hl,kn}$.

In particular $O_{ij,hl,kn}$ is singular: locally the variety $(X^{SS}(m)//G, O_{ij,hl,kn})$ is isomorphic to the toric variety

$$\mathbb{C}[T_1, T_2, T_3, T_4, T_5]/(T_1T_2T_3 - T_4T_5).$$

3. there are three indexes h, i, j s.t. $m_h + m_i + m_j = |m|/3$, then the quotient includes a point O_{hij} that correspond to the minimal, closed, strictly semi-stable orbit Gx such that $x_h = x_i = x_j$ and x_k, x_l, x_n are collinear. The point O_{hij} is non singular.

If ξ is of type 3 in Theorem 4.1, then $\theta_{\pm k}^{-1}(\xi)$ may be a point or $\mathbb{P}^3(\mathbb{C})$; in the last case $\theta_{\pm k}$ blows down $\mathbb{P}^3(\mathbb{C})$ to the point ξ :

$$\theta_{-k}^{-1}(\xi) \cong \begin{cases} \mathbb{P}^3(\mathbb{C}) & k \notin K \\ \text{point} & k \in K \end{cases} \quad \theta_{+k}^{-1}(\xi) \cong \begin{cases} \text{point} & k \notin K \\ \mathbb{P}^3(\mathbb{C}) & k \in K \end{cases} \quad (12)$$

If ξ is of type 1, then $\xi \in C_{ij} \cong \mathbb{P}^1(\mathbb{C})$ and $\theta_{\pm k}^{-1}(C_{ij})$ has dimension two (in particular it is one of the two dimensional quotients $\mathbb{P}^3(\mathbb{C})^5(m')/PSL_3(\mathbb{C})$): it is a small contraction.

As the end we can consider $\theta_{\pm k} : X^S(\widehat{m})/G \rightarrow X^{SS}(m)//G$ as the eventual composition of a blow-down map with a small map; moreover if $X^{SS}(m)//G$ is singular, then $\theta_{\pm k}$ is a resolution of singularities.

The main result of this section is the following formula for the Poincaré polynomial of an arbitrary categorical quotient $Y = X^{SS}(m)//G$:

Theorem 4.2. *Let $m = (m_1, \dots, m_6)$ be a polarization such that $0 < m_i < \frac{1}{3} \sum_i m_i, m_i \geq m_{i+1}$, then there may be five different cases:*

1. if

$$m_1 + m_2 + m_3 < \frac{2}{3}|m|,$$

$$\text{then } IP(Y) = 1 + 6t^2 + 7t^4 + 6t^6 + t^8;$$

2. if

$$\begin{cases} m_1 + m_2 + m_3 \geq \frac{2}{3}|m| \\ m_1 + m_2 + m_4 < \frac{2}{3}|m| \end{cases},$$

$$\text{then } IP(Y) = 1 + 5t^2 + 6t^4 + 5t^6 + t^8;$$

3. if

$$\begin{cases} m_1 + m_2 + m_4 \geq \frac{2}{3}|m| \\ m_1 + m_2 + m_5 < \frac{2}{3}|m| \\ m_1 + m_3 + m_4 < \frac{2}{3}|m| \end{cases},$$

$$\text{then } IP(Y) = 1 + 4t^2 + 5t^4 + 4t^6 + t^8;$$

4. if

$$\begin{cases} m_1 + m_2 + m_5 \geq \frac{2}{3}|m| \\ m_1 + m_2 + m_6 < \frac{2}{3}|m| \end{cases} \quad \text{or} \quad \begin{cases} m_1 + m_3 + m_4 \geq \frac{2}{3}|m| \\ m_2 + m_3 + m_4 < \frac{2}{3}|m| \end{cases},$$

$$\text{then } IP(Y) = 1 + 3t^2 + 4t^4 + 3t^6 + t^8;$$

5. if

$$m_1 + m_2 + m_6 \geq \frac{2}{3}|m| \quad \text{or} \quad m_2 + m_3 + m_4 \geq \frac{2}{3}|m|,$$

$$\text{then } IP(Y) = 1 + 2t^2 + 3t^4 + 2t^6 + t^8.$$

Proof. First of all let us observe that given two quotients $Y(m)$ and $Y(m')$ (geometric or categorical) it is always possible to find a finite sequence of algebraic maps $\theta_{\pm k}$ such that $Y(m) \xrightarrow{\theta_{\pm i}} \dots \xrightarrow{\theta_{\pm j}} Y(m')$, where each $\theta_{\pm k}$ is a composition of a blow-down (or blow-up) map with a small map (or only one of these).

If θ_{+k} is a small map, then it induces an isomorphism from the (intersection) homology of $Y(\widehat{m})$ to the intersection homology of $Y(m)$. It means that when we cross a wall of the G -ample cone, the intersection Betti numbers vary if the maps $\theta_{\pm k}$ involve some blow-ups and blow-downs.

As we have seen the map $\theta_{\pm k} : Y(\widehat{m}) \rightarrow Y(m)$ is a blow-down map if there is at least a point $\xi = O_{hij} \in Y(m)$ of type 3 (Theorem 4.1), that satisfies (12):

$$\theta_{\pm k}^{-1}(O_{hij}) = \theta_{\pm k}^{-1} \left(\phi(\overline{V^C}_{hij} \cup \overline{V^L}_{[6] \setminus \{h,i,j\}}) \right) = \widehat{\phi}(\overline{V^L}_{[6] \setminus \{h,i,j\}}) \cong \mathbb{P}^3(\mathbb{C}),$$

where ϕ and $\widehat{\phi}$ are the projection to the quotients $Y(m)$ and $Y(\widehat{m})$. In other words, $\theta_{\pm k}$ is a blow-down if $X^S(\widehat{m})$ contains $V^L_{[6] \setminus \{h,i,j\}}$ and not V^C_{hij} .

How many points O_{hij} there may be in a categorical quotient $Y(m)$? The answer is at most four and there are two possible sets of points:

$$\begin{matrix} O_{456}, & O_{356}, & O_{256}, & O_{156}, & \text{or} \\ O_{456}, & O_{356}, & O_{346}, & O_{345}. \end{matrix}$$

For example if $m = (666621)$ then $O_{156}, O_{256}, O_{356}, O_{456} \in X^{SS}(m)//G$ (this quotient is particularly easy to compute: it is $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$).

In this way, in order to compute the intersection Betti numbers of a quotient

$Y(m)$ (both geometric and categorical) it is sufficient to know the intersection Betti numbers for a quotient $Y(m')$ and then check how many V_K^L sets (with $|K| = 3$) are NOT included in $X^S(m)$: in fact for what we have just observed, we can always “connect” two quotients by a finite sequence of maps that change the Betti numbers if and only if they are blow-ups or blow-down of at most four copies of $\mathbb{P}^3(\mathbb{C})$. These copies of $\mathbb{P}^3(\mathbb{C})$ are detected by the sets V_K^L that are contained in $X^S(m)$, where K is one of the following:

$$456, 356, 256, 156 \quad \text{or} \quad 456, 356, 346, 345.$$

Consider now the chamber of $C^G(X)$ that contains the polarization $(5, 5, 5, 5, 1, 1)$; then for every polarization m in this chamber we have $Y(m) = \mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$ and moreover there are four sets V_K^L that are not included in $X^S(m)$. Its intersection Betti numbers are well-known and now we are able to compute all the Betti numbers:

- if m' is such that $X^S(m')$ does NOT contain four sets V_K^L then its intersection Betti numbers are the same of $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$,
- if m' is such that $X^S(m')$ does NOT contain three sets V_K^L then its intersection Betti numbers are the same of $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$ blown-up in one point,
- if m' is such that $X^S(m')$ does NOT contain two sets V_K^L then its intersection Betti numbers are the same of $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$ blown-up in two points,
- if m' is such that $X^S(m')$ does NOT contain one sets V_K^L then its intersection Betti numbers are the same of $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$ blown-up in three points,
- if m' is such that $X^S(m')$ contains all sets V_K^L then its intersection Betti numbers are the same of $\mathbb{P}^2(\mathbb{C}) \times \mathbb{P}^2(\mathbb{C})$ blown-up in four points.

The relations of Theorem 4.2 compute the number of V_K^L sets that are included in $X^S(m)$. ■

References

- [1] I.V. Dolgachev, *Lectures on Invariant Theory*, Cambridge University Press, Lecture Note Series **296**, 2003.
- [2] I.V. Dolgachev, Y. Hu, *Variations of geometric invariant theory quotients*, Duke Math. Jour., **68**, 1992, 151-184.
- [3] J.-C. Hausmann, A. Knutson, *The cohomology ring of polygon spaces*, Ann. Inst. Fourier (Grenoble), **48**, 1998, no. 1, 281-321.

- [4] Y. Hu, *The geometry and topology of quotient varieties of torus actions*, Publ. Math. IHES, **87**, 1998, 5-51.
- [5] F. Incensi, *GIT quotients of products of projective planes*, Accepted by Rendiconti Seminari Matematici dell'Università di Padova, 2009.
- [6] D. Mumford, J. Fogarty, F. Kirwan, *Geometric Invariant Theory*, Springer-Verlag, third edition, 1994.
- [7] M. Thaddeus, *Geometric invariant theory and flips*, Jour. Amer. Math. Soc., **9**, 1996, 691-723.

FRANCESCA INCENSI

Dipartimento di Matematica, Università di Bologna, Italy

E-mail address: *incensi@dm.unibo.it*